

Crossover Phase Diagram of 2D Heisenberg Ferro- and Antiferromagnets

Alexander Sokol

*Department of Physics, University of Illinois at Urbana-Champaign, Urbana, IL 61801-3080
and L.D. Landau Institute, Moscow, Russia*

Norbert Elstner

Service de Physique Théorique, CEA-Saclay, 91191 Gif-sur-Yvette Cedex, France

Rajiv R.P. Singh

*Department of Physics, University of California, Davis, CA 95616
(cond-mat/9505148, May 29, 1995)*

We propose a single crossover phase diagram applicable to 2D collinear Heisenberg antiferromagnets (AFMs) and ferromagnets (FMs), and show that the scaling regimes of AFMs and FMs are in one-to-one correspondence. The phase diagram is split into classical and quantum regions. Our two key results are: (i) in the classical region, the AFMs and FMs exhibit nearly identical behavior near their respective ordering wavevectors, which we observe for $S=1$ and higher using series expansions; and (ii) in the quantum region, quantum critical (QC) regime is present not only for the AFMs, but for FMs as well.

I. INTRODUCTION

We report a study of two-dimensional, collinear, spin- S antiferromagnets (AFMs) and ferromagnets (FMs) aimed at better understanding the magnetic behavior of both, and in particular the role of quantum versus classical fluctuations in these systems. A number of quasi-2D experimental systems, including spin- $\frac{1}{2}$ AFMs La_2CuO_4 and $\text{Sr}_2\text{CuO}_2\text{Cl}_2$ [1,2], spin-1 AFMs La_2NiO_4 and K_2NiF_4 [3], spin- $\frac{5}{2}$ AFM Rb_2MnF_4 [4] as well as spin- $\frac{1}{2}$ FMs such as K_2CuF_4 [5] are well described over a range of temperatures by the 2D Heisenberg model

$$H = J \sum_{\langle ij \rangle} \mathbf{S}_i \mathbf{S}_j \quad (1)$$

on a square lattice, which we study by high temperature series expansion methods. We expect much of our results to apply to other lattices having collinear long-range order at $T = 0$.

Let us begin by considering the relevant energy scales in the problem. One important energy scale for both AFMs ($J > 0$) and FMs ($J < 0$) is the $T = 0$ spin stiffness ρ_s , defined as rigidity with respect to a twist in the magnetic structure. Quantum $1/S$ corrections make this quantity different for AFMs and FMs with the same value of spin; for model (1), the order of magnitude is however the same, $\rho_s^{\text{AFM}} \sim JS^2$, $\rho_s^{\text{FM}} = JS^2$ (note that the FM value is exact).

As shown by Chakravarty, Halperin, and Nelson [6] for AFMs, and by Kopietz and Chakravarty [7] for FMs, the asymptotic $T \rightarrow 0$ magnetic behavior for these models obeys scaling and depends on only two dimensionful parameters: the energy scale ρ_s , and a quantity which sets the overall length scale and can be obtained from the $q \rightarrow 0$ limit of the spin wave dispersion $\epsilon(q)$. For AFMs, $\epsilon(q)$ is linear and one defines the $T = 0$ spin wave

velocity as $c = \lim_{q \rightarrow 0} \epsilon(q)/q \sim JSa$. For FMs, ϵ_q is quadratic and one defines the $T = 0$ spin wave stiffness $\rho = \lim_{q \rightarrow 0} \epsilon(q)/q^2 \sim JSa^2$. Note that the spin wave stiffness ρ has the dimension of energy $\times$ length², while the spin stiffness ρ_s has the dimension of energy.

For the Heisenberg model, the spin wave spectrum is well-defined for all q and its upper bound, reached at the boundary of the Brillouin zone where $q \sim 1/a$, can be estimated as $\Lambda \sim c/a \sim JS$ for the AFMs, or $\Lambda \sim \rho/a^2 \sim JS$ for the FMs, i.e. not only ρ_s , but also $\Lambda \sim JS$ has the same order of magnitude in AFMs and FMs.

Whereas the $T \rightarrow 0$ behavior depends on ρ_s and c for AFMs, or ρ_s and ρ for FMs, the $T \rightarrow \infty$ behavior (the Curie-Weiss law) depends on $\rho_{\text{CL}} = JS(S+1)$ and the lattice constant a . In what follows, we show that this simple observation is part of a general picture where the low temperature behavior, which may contain several scaling regimes, at $T \sim \Lambda$ crosses over to the high-temperature behavior, which may also contain several scaling regimes. A similar crossover in 3D occurs *below* the long range ordering temperature (which is zero in 2D), and was studied by Vaks, Larkin, and Pikin [8].

In model (1), the ratio $\Lambda/\rho_s \sim 1/S$ can be made arbitrarily small by increasing S , but it cannot be made arbitrarily large as $S \geq \frac{1}{2}$. Nevertheless, the case $\Lambda/\rho_s \gg 1$ is of great interest. First, there exist models where ρ_s goes to zero while Λ does not, in which case $\Lambda/\rho_s \gg 1$ applies rigorously; an example of such a system is the two-layer Heisenberg model [9,10] near the critical point where the Néel AFM long-range order vanishes. Second, for the $S = 1/2$ AFM model (1), the ratio $\Lambda/\rho_s \sim 10 \gg 1$ is quite large; furthermore, there exists evidence [11,12,9,13,10], first pointed out by Chubukov and Sachdev, that at intermediate temperatures this model is indeed in the quantum critical limit $\Lambda/\rho_s \gg 1$.




FIG. 1. A phase diagram of spin- S quantum antiferromagnets (AFMs) and ferromagnets (FMs); see Table I for the corresponding scaling and crossover expressions for the correlation length. All regime boundaries are gradual crossovers rather than phase transitions, and their positions can only be defined within numerical factors of order unity. The behavior in the $T/\Lambda \gg 1$ region is in the universality class of 2D classical magnets, where magnetic properties of AFMs and FMs for the same value of S are the same near their respective ordering wavevectors, and depend on T/ρ_{CL} , where $\rho_{\text{CL}} = JS(S+1)$. The classical behavior includes Curie-Weiss regime ($T \gg \max(\rho_s, \Lambda)$) and classical scaling regime ($\Lambda \ll T \ll \rho_{\text{CL}}$), which are separated by a fairly wide *classical crossover* regime for $\Lambda \ll T \sim \rho_{\text{CL}}$, where scaling does not hold, but nevertheless $\xi_{\text{AFM}} \approx \xi_{\text{FM}} \approx a\psi_{\text{CL}}(T/JS(S+1))$. In the region $T/\Lambda \ll 1$ quantum effects are important and therefore AFMs and FMs behave differently. For the AFMs, this region is in the universality class of the QNL σ model. Here these scaling regimes are found for both AFMs and FMs: renormalized classical (RC) regime for $T \ll \min(\rho_s, \Lambda)$; quantum critical (QC) regime for $\rho_s \ll T \ll \Lambda$; and a wide quantum crossover regime for $T \sim \rho_s \ll \Lambda$, where the correlation length remains a universal function (different for AFMs and FMs) of the thermal de Broglie wavelength for spin waves ($\lambda_T = c/T$ or $\lambda_T = (\rho/T)^{1/2}$) and the ratio T/ρ_s .

Regime	Correlation Length
Curie-Weiss $T \gg \max(\rho_{\text{CL}}, \Lambda)$	$\xi_{\text{AFM}} = \xi_{\text{FM}} \lesssim a$ Properties depend on T/ρ_{CL}
Classical Crossover $T \sim \rho_{\text{CL}} \gg \Lambda$	$\xi_{\text{AFM}} = \xi_{\text{FM}} = a\psi_{\text{CL}}(T/\rho_{\text{CL}})$
Classical $\rho_{\text{CL}} \gg T \gg \Lambda$	$\xi_{\text{AFM}} = \xi_{\text{FM}} \sim a(T/\rho_{\text{CL}}) \times \exp(2\pi\rho_{\text{CL}}/T)$
Quantum Critical $\rho_s \ll T \ll \Lambda$	$\xi_{\text{AFM}} \sim c/T,$ $\xi_{\text{FM}} \sim (\rho/T \log(T/\rho_s))^{1/2}$
Quantum Crossover $T \sim \rho_s \ll \Lambda$	$\xi_{\text{AFM}} = (c/T) \phi_{\text{AFM}}(T/\rho_s),$ $\xi_{\text{FM}} = (\rho/T)^{1/2} \phi_{\text{FM}}(T/\rho_s)$
Renormalized Classical $T \ll \min(\rho_s, \Lambda)$	$\xi_{\text{AFM}} \sim (c/\rho_s) \exp(2\pi\rho_s/T),$ $\xi_{\text{FM}} \sim (\rho T/\rho_s)^{1/2} \exp(2\pi\rho_s/T)$

TABLE I. Correlation length in the scaling and crossover regimes shown in the phase diagram of Fig.1, after Refs. [14,6,7,11], and this work. The sign $\sim$ indicates the presence of a numerical prefactor. The functions ϕ_{AFM} and ϕ_{FM} are universal; the function ψ_{CL} is not, for instance, it explicitly depends on the ratio of the next-nearest and nearest-neighbor exchange couplings.

II. CURIE-WEISS REGIME

At high temperatures $T \gg \max(\rho_s, \Lambda)$, the spins are weakly correlated and one can keep only a few leading terms in the high temperature series expansion. For model (1), this temperature range corresponds to $T \gg \max(JS, JS^2)$. The corresponding mean-field theory yields the familiar Curie-Weiss law.

III. RENORMALIZED CLASSICAL REGIME

In the opposite limit of $T \ll \min(\rho_s, \Lambda)$, the behavior is of the renormalized classical (RC) type. This regime was studied in detail in Refs. [6,7,11], where magnetic properties were calculated by mapping the system onto the classical nonlinear σ -model and setting the momentum-integration cutoff to be proportional to the thermal de Broglie wave vector q_T , defined such that $\epsilon(q_T) \sim T$. The predicted $\xi(T)$ has the same form for AFMs ($J > 0$) and FMs ($J < 0$), when expressed in terms of their respective thermal de Broglie wavelengths $\lambda_T = 1/q_T$:

$$\xi_{\text{RC}} = \frac{e}{8} \lambda_T \frac{T}{2\pi\rho_s} \exp\left(\frac{2\pi\rho_s}{T}\right), \quad (2)$$

(here the exact value of the prefactor is obtained from the correlation length of the classical $O(3)$ nonlinear- σ model in the minimal subtraction scheme [15,16]). However, not only the values of ρ_s differ for FMs and AFMs, but also λ_T has qualitatively different temperature dependences:

$$\lambda_T = \begin{cases} c/T & \text{for AFMs} \\ \sqrt{\rho/T} & \text{for FMs} \end{cases}. \quad (3)$$

An important question is upto what temperatures should this RC expression hold? We recall that when T is larger than the upper bound of the $T = 0$ spin wave spectrum, Λ , there can be no spin waves with wavelengths that are comparable to the thermal de Broglie wavelength. Since such spin waves are important for RC theory, one would expect it to be applicable only for $T \ll \min(\Lambda, \rho_s)$, which for model (1) translates into $T \sim \min(JS, JS^2)$ (here, we ignore numerical factors of order unity). At higher temperatures new physics must arise.

The Curie-Weiss regime occurs for $T \gg \max(\rho_s, \Lambda)$ and the RC regime for $T \ll \min(\rho_s, \Lambda)$. These two regimes exist for all S . We now turn to the discussion of the regimes where the temperature is larger than one of the scales ρ_s or Λ , but smaller than the other.




FIG. 2. A semi-log plot of ξ/a vs. $T/(JS(S+1))$ for AFMs (solid symbols) and FMs (open symbols) for $S = 1/2$ (circles) and $S = 5/2$ (diamonds). We also plot the data from Ref. [14] for the classical $S \rightarrow \infty$ model (solid line). For $S = 1/2$, AFM and FM correlation lengths are different and neither agrees with the $S \rightarrow \infty$ limit. For $S = 5/2$, AFM and FM correlation lengths agree *both* with each other and with the $S \rightarrow \infty$ model. The data for $S = 5/2$ is plotted as representative for large spin models; for all studied spins $S > 1$, the behavior is similar although the agreement between ξ_{AFM} and ξ_{FM} for $S = 3/2$ and $S = 2$ is not as striking as for $S = 5/2$. This data collapse provides evidence that any quantum effects, which are expected to be different for AFMs and FMs, become unimportant already for fairly small spins $S > 1$, in agreement with the conjecture due to Ref. [17].

IV. CLASSICAL AND CLASSICAL CROSSOVER REGIMES

In our earlier work in collaboration with Greven and Birgeneau [17], we proposed a scaling crossover scenario to describe substantial deviations from RC behavior observed in the neutron scattering experiments for AFMs with spin-one and larger. This scenario calls for a crossover to the behavior of the $S \rightarrow \infty$ classical system when spin wave energies for all wavevectors in the Brillouin zone become smaller than the temperature, i.e. for $T \gtrsim \Lambda$. The correlation length at this RC to classical boundary, obtained from Eqs.(2), is exponentially large for large spin,

$$\left. \frac{\xi}{a} \right|_{T \sim \Lambda} \sim \frac{\exp(S)}{S} \gg 1 \quad \text{for } S \gg 1. \quad (4)$$

In [17], the arguments in favor of RC to classical crossover were based on data collapse for large spin when plotted versus $T/(JS(S+1))$. The studied values of spin were small enough ($S = 1/2$ to $S = 5/2$) to make the results sensitive to the choice of the renormalized temperature as $T/(JS(S+1))$ or T/JS^2 .

Here, we resolve this arbitrariness by studying antiferro- and ferromagnets simultaneously. AFMs and

FMs have the same classical limit, but the quantum effects differ and therefore the difference between AFMs and FMs can be used as a probe for the strength of quantum effects. It turns out that for any $S > 1$, in the temperature range of interest ($\xi \sim 10^1$), the correlation length calculated by high temperature expansions is nearly the same for AFMs and FMs with the same value of S (Fig.2). Therefore, it must be determined by classical rather than quantum physics. We emphasize that such an argument does not rely on any particular way of collapsing the data, because it is drawn from comparisons of AFM and FM models for the same spin. On the other hand, for $S = 1/2$ the difference between ξ_{AFM} and ξ_{FM} is always large and temperature-dependent, which indicates the importance of quantum physics for this value of S . Spin-one appears to be a borderline case.

Having established that in the temperature range $T \gg \Lambda$ finite-spin models behave similarly to the $S \rightarrow \infty$ classical magnet, we now comment on the behavior of the latter. The asymptotic low-temperature behavior of the classical 2D Heisenberg antiferromagnet is given by [16]:

$$\frac{\xi_{\text{CL}}}{a} = \frac{e}{8} \frac{e^{-\pi/2}}{\sqrt{32}} \frac{T}{2\pi\rho_{\text{CL}}} \exp\left(\frac{2\pi\rho_{\text{CL}}}{T}\right), \quad (5)$$

where $\rho_{\text{CL}} = JS(S+1)$. It turns out that because of the very small numerical value of the prefactor (~ 0.01), this formula becomes accurate only for very low temperatures, $T/\rho_{\text{CL}} \lesssim 0.6$ ($\xi \gtrsim 100$). We label the intermediate temperature range where neither Curie-Weiss nor classical scaling behavior apply the *classical crossover* regime. In this regime, neither large nor small-temperature asymptotic expressions describe the correlations accurately, nevertheless, all properties of the model are nearly the same for AFMs and FMs, and depend on T through T/ρ_{CL} only.

V. QUANTUM CRITICAL AND QUANTUM CROSSOVER REGIMES

We call *quantum critical* that behavior which can formally be obtained for $\rho_s \ll T \ll \Lambda$. For the antiferromagnets, this regime is the asymptotic *quantum critical* (QC) regime [6,11]. The correlation length ξ is proportional to $\lambda_T = c/T$ with corrections that depend universally on T/ρ_s , and to leading order the dominant frequency scale for spin fluctuations is $\bar{\omega} \sim T$.

For the FMs the spin wave spectrum is quadratic, which causes infrared ($q \rightarrow 0$) log divergences for $\rho_s \ll T \ll \Lambda$. Such divergences (not present for AFMs) are cutoff by ρ_s , and lead to the following multiplicative correction to the correlation length:

$$\xi_{\text{FM}} \sim (\rho/T)^{1/2} \log^{-1/2}(T/\rho_s), \quad (6)$$

in either the sigma model or the Schwinger boson formalism.

Note that such a behavior cannot be the true quantum critical behavior because the limit $\rho_s \rightarrow 0$ leads to singular ξ at finite T . Therefore, Eq.(6) must fail as $\rho_s \rightarrow 0$, and the $T = 0$ quantum ferromagnet-paramagnet phase transition is likely to belong to a different universality class [18]. Nevertheless, lacking a better name and for compatibility with the AFM nomenclature, we call this behavior a *ferromagnetic quantum critical* regime.

The applicability of the QC description to model (1) for $S = 1/2$ AFM is widely discussed in the literature [2,11,12,9,13]. A similar analysis for the ferromagnets requires detailed calculations of their properties in the QC regime, and is beyond the framework of this paper.

VI. CONCLUSION

In this paper, we present a complete phase diagram of spin- S antiferromagnets (AFMs) and ferromagnets (FMs). We show that comparing AFMs and FMs allows one to elucidate crossover effects from the quantum regimes (including the renormalized classical regime where, despite its name, quantum fluctuations are essential) to the purely classical regimes, where AFMs and FMs exhibit identical behavior near their respective ordering wavevectors. For a detailed description of the phase diagram and the proposed new regimes, we refer the reader to Fig.1 and Table 1.

For the antiferromagnets, both our series expansion calculations and the neutron data [2,17] suggest that the regime boundaries in Fig.1 are positioned such that the universal $T \ll \Lambda$ scaling theory, free of any lattice corrections, is not obeyed for spin-one and larger at any numerically or experimentally accessible values of the correlation length. The data also highlights the disagreement in the magnitude of spin correlations even for $S = 1/2$ at all accessible temperatures with the existing RC predictions, whereas $\xi(T)$ was found [2] to agree remarkably well with the RC theory.

Our results suggest that an accurate analytical theory for $S \geq 1$ in the experimentally relevant temperature range can be constructed by taking into account leading quantum corrections about the $S = \infty$ limit on a lattice. The important step here would be to obtain an analytic approximation which is valid in the *classical crossover regime*. On the other hand, any theory based on a purely continuum description, such as the QNL σ model without lattice corrections, is clearly inadequate for any spin larger than $S = 1/2$.

We are grateful to R.J. Birgeneau, S. Chakravarty, A.V. Chubukov, M. Greven, Th. Jolicœur, D. Pines, S. Sachdev, and S. Sondhi for many useful discussions. This work is supported by the NSF Grant No. DMR 93-18537. One of us (A.S.) is supported by an Alfred P. Sloan Research Fellowship.

-
- [1] B. Keimer *et al.*, Phys. Rev. B **46**, 14034 (1992); R.J. Birgeneau *et al.* (unpublished)
 - [2] M. Greven *et al.*, Phys. Rev. Lett. **72**, 1096 (1994), and Z. Phys. B **96**, 465 (1995)
 - [3] K. Nakajima *et al.*, Z. Phys. B **96**, 479 (1995); R.J. Birgeneau, Phys. Rev. B **41**, 2514 (1990), and references therein.
 - [4] R.A. Cowley *et al.*, Phys. Rev. B **15**, 4292 (1977).
 - [5] L. J. Dejongh and A. R. Midema, Advances in Physics, **23**, 1 (1994).
 - [6] S. Chakravarty, B.I. Halperin, and D.R. Nelson, Phys. Rev. B **39**, 2344 (1989). The exact value of the prefactor in the expression for $\xi(T)$ has been calculated in Ref. [15].
 - [7] P. Kopietz and S. Chakravarty, Phys. Rev. B **40**, 4858 (1989).
 - [8] V.G. Vaks, A.I. Larkin, and S.A. Pikin, Sov. Phys. JETP **26**, 188 (1968).
 - [9] A.W. Sandvik and D.J. Scalapino, Phys. Rev. Lett. **72**, 2777 (1993); A.W. Sandvik and A.V. Chubukov, unpublished. A.J. Millis and H. Monien [Phys. Rev. Lett. **70**, 2810 (1994)] proposed to use the two-layer Heisenberg model to describe bilayer interaction in YBCO materials.
 - [10] A.W. Sandvik, A.V. Chubukov, and S. Sachdev, preprint (1995). Available as cond-mat/9502012.
 - [11] A.V. Chubukov, S. Sachdev, and J. Ye, Phys. Rev. B **49**, 11919 (1994); A.V. Chubukov and S. Sachdev, Phys. Rev. Lett. **71**, 169 (1993).
 - [12] A. Sokol, R.L. Glenister, and R.R.P. Singh, Phys. Rev. Lett., **72**, 1549 (1994).
 - [13] N. Elstner, R. L. Glenister, R. R. P. Singh, and A. Sokol, Phys. Rev. B **51**, 8984 (1995).
 - [14] M. Lüscher and P. Weisz, Nucl. Phys. B **300**, 325 (1988); S.H. Shenker and J. Tobochnik, Phys. Rev. B **22**, 4462 (1980); D.N. Lambeth and H.E. Stanley, Phys. Rev. B **12**, 5302 (1975).
 - [15] P. Hasenfratz M. Maggiore and F. Niedermayer, Phys. Lett. B **245**, 522 (1990); P. Hasenfratz and F. Niedermayer, Phys. Lett. B **268**, 231 (1991).
 - [16] Th. Jolicœur and J.C. LeGuillou, Mod. Phys. Lett. B **5**, 593 (1991).
 - [17] N. Elstner, A. Sokol, R.R.P. Singh, M. Greven, R.J. Birgeneau, preprint (1995).
 - [18] S. Sachdev, Z. Phys. B **94**, 469 (1994)